

Pre and Post Photofacial

Redness, Discoloration, Sun Damage, Facial Veins and Rosacea

Pre-Treatment:

- Avoid indoor/outdoor tanning, self tanners and spray tanning for 4 weeks prior.
- Discontinue the use of Trentinoin, Retin-A, Retinol, Renova and any products containing AHA's and BHA's 2 weeks prior to treatment.
- Review medications with esthetician to check photo sensitivity.

Contraindications- Herpes 1 or 2 within treatment area, diabetes needs to be controlled, history of heart urticarial as you may develop hives, if on accutane or within last 6 months, if you take anti coagulants or blood thinners-needs doctor approval, history of vitiligo, eczema, psoriasis, allergic dermatitis, autoimmune disease, collagen affecting diseases and Ehlers-Danios Syndrome.

CAN NOT HAVE TREATMENT

- **Slow healing**
- **Skin Cancer**
- **Active infections/Immunosuppression**
- **Open Lesions**
- **Pregnancy**
- **Blood Disorders**
- **Keloid Scarring**
- **Use of Steroids**

Post-Treatment

- Skin may appear red or blotchy in the area for 24-72 hours post treatment.
- Wash area with mild cleanser.
- Apply post cream/ointment twice daily.
- Take Tylenol as needed for discomfort.
- No shaving in area that is red and swollen.
- Avoid sun exposure, tanning beds, self tanners and spray tanning for the rest of treatments.
- Apply SPF daily 30+ for the rest of treatment.
- Avoid alcohol, spicy foods, caffeine and workouts for 24-48 hours post treatment.